

68677

43.1.-2015.

ISSN 1027-4634

# МАТЕМАТИЧНІ СТУДІЇ

---

МАТЕМАТИЧНИ  
STUDII

Том 43, No. 1    **2015**    Vol. 43, No. 1

## МАТЕМАТИЧНІ СТУДІЇ

Праці Львівського математичного товариства

ISSN 1027-4634

Журнал присвячено дослідженням в усіх областях математики. Приймаються оригінальні статті середнього розміру; виняток можливий для оглядових статей. Статті публікуються англійською, німецькою, російською, українською та французькою мовами. Виходить щоквартально.

Видається видавництвом «ВНТЛ–Класика» для Львівського математичного товариства за фінансової та організаційної підтримки Львівського національного університету.

### Редактори:

М.М.Зарічний (Львів), О.Б.Скасків (Львів),  
М.М.Шеремета (Львів)

### Редакційна колегія:

О.Д.Артемович (Івано-Франківськ; Краків, Польща), Т.О.Банакх (Львів), М.М.Бокало (Львів), Ю.Д.Головатий (Львів), Р.І.Григорчук (Москва, Росія), Р.О.Гринів (Львів), І.Й.Гуран (Львів), А.А.Дороговцев (Київ), Ю.А.Дрозд (Київ), С.Д.Івасишен (Київ), В.В.Кириченко (Київ), М.Я.Комарницький (Львів), Б.І.Копитко (Львів), А.А.Кореновський (Одеса), Р.Коті (Париж, Франція), О.В.Лопушанський (Львів; Жешув, Польща), Я.В.Микитюк (Львів), Й.В.Островський (Харків; Анкара, Туреччина), О.А.Панков (Балтімор, США), М.М.Попов (Чернівці), І.В.Протасов (Київ), Б.Й.Пташник (Львів), Д.Реповш (Любляна, Словенія), В.Г.Самойленко (Київ), А.М.Седлецький (Москва, Росія), О.Г.Сторож (Львів), В.І.Сушанський (Глівіце, Польща), С.Ю.Фаворов (Харків), Б.Н.Хабібуллін (Уфа, Росія), І.Е.Чижиков (Львів)

### Технічний редактор:

Д.Ю.Зікрач

### Адреса редакції:

Україна, 79000 Львів,  
Університетська, 1  
Львівський національний університет,  
механіко-математичний факультет,  
Математичні студії  
e-mail: matstud@franko.lviv.ua

### Інформація про передплату:

З питань передплати звертатись за адресою:  
ВНТЛ–Класика  
а/с 10249, Львів, 79006  
e-mail: info@vntl.com, <http://www.vntl.com>

## MATEMATYCHNI STUDIИ

Proceedings of the Lviv Mathematical Society

ISSN 1027-4634

Journal is devoted to research in all fields of mathematics. Original papers of moderate length are accepted; exception is possible for survey articles. Languages accepted are: English, French, German, Russian, and Ukrainian. Published quarterly.

Published for the Lviv Mathematical Society by VNTL Publishers with financial and organizing support of the Lviv National University.

### Editors-in-Chief:

M.M.Sheremeta (Lviv), O.B.Skaskiv (Lviv),  
M.M.Zarichnyi (Lviv)

### Editorial Board:

O.D.Artemovych (Ivano-Frankivs'k; Kraków, Poland), T.O.Banakh (Lviv), M.M.Bokalo (Lviv), R.Cauty (Paris, France), I.E.Chyzykhov (Lviv), A.A.Dorogovtsev (Kyiv), Yu.A.Drozd (Kyiv), S.Yu.Favorov (Kharkiv), Yu.D.Golovaty (Lviv), R.I.Grigorchuk (Moscow, Russia), I.Yo.Guran (Lviv), R.O.Hryniv (Lviv), S.D.Ivasyshen (Kyiv), B.N.Khabibullin (Ufa, Russia), V.V.Kirichenko (Kyiv), M.Ya.Komarnytskyi (Lviv), B.I.Kopytko (Lviv), A.A.Korenovskii (Odessa), O.V.Lopushanskyi (Lviv; Rzeszów, Poland), Ya.V.Mykytyuk (Lviv), I.V.Ostrovskii (Kharkiv; Ankara, Turkey), O.A.Pankov (Baltimore, USA), M.M.Popov (Chernivtsi), I.V.Protasov (Kyiv), B.Yo.Ptashnyk (Lviv), D.Repovš (Ljubljana, Slovenia), V.G.Samoilenko (Kyiv), A.M.Sedletskii (Moscow, Russia), O.G.Storozh (Lviv), V.I.Sushchansky (Glivice, Poland)

### Technical editor:

D.Yu.Zikrach

### Address:

Matematychni Studii  
Dept. of Mechanics and Mathematics,  
Lviv National University,  
1 Universytetska St.,  
79000, Lviv, Ukraine  
e-mail: matstud@franko.lviv.ua

### Subscription information:

Subscription information can be obtained by the address:  
VNTL Publishers  
P.O.Box 10249, Lviv, 79006, Ukraine  
e-mail: info@vntl.com, <http://www.vntl.com>

## З М І С Т

ZHUCHOK YUL. V. Free $n$ -nilpotent trioids.....	3
LUKASHENKO M. P. Derivations as homomorphisms or anti-homomorphisms in differentially semiprime rings.....	12
KURYLIAK A. O., SHAPOVALOVSKA L. O. Wiman's inequality for entire functions of several complex variables with rapidly oscillating coefficients.....	16
HECTEPENKO B. B. Сукупні властивості відображень, які пов'язані із замкненістю графіка .....	27
KAPLOVA O. Сильно нарізно неперервні функції і одна характеристика відкритих множин в ящиківому добутку .....	36
FAVOROV S. YU., GIRYA N. P. On a property of pairs of almost periodic zero sets.....	43
TARGONSKIY A. L. Деякі нерівності для внутрішніх радіусів попарно- неперетинних областей та відкритих множин .....	51
KHOROSHCHAK V. S., KONDRATYUK A. A. The Riesz measures and a repre- sentation of multiplicatively periodic $\delta$ -subharmonic functions in a punc- tured Euclidean space .....	61
DEREV'YANKO T. O., KYRYLICH V. M. Оптимальне керування квазіліній- ною гіперболічною системою, що описує попит Слуцького.....	66
MASLYUCHENKO V. K., MASLYUCHENKO O. V., MYRONYK O. D. On $L$ -separatedness and $L$ -regularity of the Ceder products.....	78

## КОРОТКІ ПОВІДОМЛЕННЯ

СИМОТЮК М. М. Метричні оцінки визначника інтерполяційної задачі, один з вузлів якої є кратним, для лінійних рівнянь з частинними похі- дними.....	88
<span style="border: 1px solid black; padding: 2px;">Гришин А. Ф.</span> , Куиць Н. В. Предельные множества Азарина для мер Радона. I.....	94

## ПРОБЛЕМИ

IL'INSKII A. I. On the division of a characteristic function by the Blaschke product .....	100
BANDURA A. I., SKASKIV O. B. Open problems for entire functions of bounded index in direction .....	103
SAVCHENKO A., ZARICHNYI M. Open problems in the theory of fuzzy metric spaces.....	110

## CONTENTS

ZHUCHOK YUL. V. Free $n$ -nilpotent trioids.....	3
LUKASHENKO M. P. Derivations as homomorphisms or anti-homomorphisms in differentially semiprime rings.....	12
KURYLIAK A. O., SHAPOVALOVSKA L. O. Wiman's inequality for entire functions of several complex variables with rapidly oscillating coefficients.....	16
NESTERENKO V. V. Aggregate properties of mappings that are associated with the closedness of a graph.....	27
KARLOVA O. Strongly separately continuous functions and a characterization of open sets in a box-product.....	36
FAVOROV S. YU., GIRYA N. P. On a property of pairs of almost periodic zero sets.....	43
TARGONSKII A. L. Some inequalities for inner radii of pair-wise disjoint domains and open sets.....	51
KHOROSHCHAK V. S., KONDRATYUK A. A. The Riesz measures and a repre- sentation of multiplicatively periodic $\delta$ -subharmonic functions in a punc- tured Euclidean space.....	61
DEREVIANKO T. O., KYRYLYCH V. M. Optimal control problem for quasili- near hyperbolic system: the Slutsky equation.....	66
MASLYUCHENKO V. K., MASLYUCHENKO O. V., MYRONYK O. D. On $L$ -separatedness and $L$ -regularity of the Ceder products.....	78

## RESEARCH ANNOUNCEMENTS

SYMOTYUK M. M. Metric estimates of the characterictic determinant of an interpolation problem with nodes, one of which is multiple, for a linear partial differential equation.....	88
<span style="border: 1px solid black; padding: 2px;">GRISHIN A. F.</span> , QUYNH N. V. Azarin limit sets for Radon measures. I.....	94

## PROBLEM SECTION

IL'INSKII A. I. On the division of a characteristic function by the Blaschke product.....	100
BANDURA A. I., SKASKIV O. B. Open problems for entire functions of bounded index in direction.....	103
SAVCHENKO A., ZARICHNYI M. Open problems in the theory of fuzzy metric spaces.....	110

УДК 512.579

YUL. V. ZHUCHOK

**FREE  $n$ -NILPOTENT TRIOIDS**Yul. V. Zhuchok. *Free  $n$ -nilpotent trioids*, Mat. Stud. **43** (2015), 3–11.

We introduce the notion of a nilpotent trioid, construct a free  $n$ -nilpotent trioid and describe its structure. We also characterize the least  $n$ -nilpotent congruence on a free trioid and give examples of nilpotent trioids of nilpotency index 2.

Юл. В. Жучок. *Свободные  $n$ -нильпотентные триоиды* // Мат. Студії. – 2015. – Т.43, №1. – С.3–11.

Введено понятие нильпотентного триоида, построен свободный  $n$ -нильпотентный триоид и описана его структура. Также охарактеризована наименьшая  $n$ -нильпотентная конгруэнция на свободном триоиде и приведены примеры нильпотентных триоидов индекса нильпотентности 2.

**1. Introduction.** J.-L. Loday and M. O. Ronco ([1]) introduced a type of algebras, called trioids, which are sets endowed with three binary associative operations  $\dashv$ ,  $\vdash$  and  $\perp$  satisfying eight axioms:  $(x \dashv y) \dashv z = x \dashv (y \vdash z)$  (T1),  $(x \vdash y) \dashv z = x \vdash (y \dashv z)$  (T2),  $(x \dashv y) \vdash z = x \vdash (y \vdash z)$  (T3),  $(x \dashv y) \dashv z = x \dashv (y \perp z)$  (T4),  $(x \perp y) \dashv z = x \perp (y \dashv z)$  (T5),  $(x \dashv y) \perp z = x \perp (y \vdash z)$  (T6),  $(x \vdash y) \perp z = x \vdash (y \perp z)$  (T7),  $(x \perp y) \vdash z = x \vdash (y \vdash z)$  (T8). Trioids have applications in the theory of trialgebras ([1]). Recall that trialgebras are linear analogs of trioids. This kind of algebras is closely related to ternary planar trees. It is well known that dialgebras (dimonoids) ([2, 3]) can be obtained from trialgebras (trioids). In the survey paper [4], numerous examples of trioids were presented. The problem of constructing free trioids was solved in [1, 4]. Free rectangular trioids were given in [5].

Nilpotency in different algebras has been extensively studied by many authors. So, the notion of a nilpotent semigroup was introduced by A. I. Malcev ([6]) and independently by B. H. Neuman and T. Taylor ([7]). The relationships between nilpotent semigroups and semigroup algebras were studied by E. Jespers and J. Okninski ([8]). Nilpotency in rings was considered in [9]. Papers [10, 11] are devoted to studying (di)nilpotent dimonoids.

This paper develops the variety theory of trioids. In Section 2 constructions of a free trioid and some other algebras are given. In Section 3 we introduce the notion of a nilpotent trioid, give examples of nilpotent trioids of nilpotency index 2 and construct a free  $n$ -nilpotent trioid. In Section 4 we introduce the notion of a 0-triband of subtrioids and in terms of 0-tribands of subtrioids describe the structure of free  $n$ -nilpotent trioids. In the final section the least  $n$ -nilpotent congruence on a free trioid is characterized.

**2. Preliminaries.** Consider free trioids (see [4]).

2010 *Mathematics Subject Classification*: 08A05, 17A30, 17D99, 20M10, 20M50.

*Keywords*: trioid; nilpotent trioid; free  $n$ -nilpotent trioid; dimonoid; semigroup; congruence.

doi:10.15330/ms.43.1.3-11

Let  $Y$  be an arbitrary nonempty set,  $\bar{Y} = \{\bar{x} \mid x \in Y\}$ ,  $X = Y \cup \bar{Y}$  and  $F[X]$  be the free semigroup on  $X$ . Let further  $P \subset F[X]$  be a subsemigroup which contains words  $w$  with the element  $\bar{x}$  ( $x \in Y$ ) occurring in  $w$  at least one time. It is easy to see that  $F[X]$  is a band of semigroups  $P$  and  $F[X] \setminus P$  [12].

Let  $w \in P$ . Denote by  $\tilde{w}$  the word obtained from  $w$  by the replacement of all letters  $\bar{x}$  ( $x \in Y$ ) with  $x$ . For instance, if  $w = x\bar{x}\bar{y}x\bar{z}$ , then  $\tilde{w} = xxyxz$ . Obviously,  $\tilde{w} \in F[X] \setminus P$ .

Define operations  $\dashv$ ,  $\vdash$  and  $\perp$  on  $P$  by

$$w \dashv u = w\tilde{u}, \quad w \vdash u = \tilde{w}u, \quad w \perp u = wu$$

for all  $w, u \in P$ . Denote the algebra  $(P, \dashv, \vdash, \perp)$  by  $\text{Frt}(Y)$ .

The proof of the following statement is similar to the proof of Proposition 1.9 from [1] obtained for the free trioid of rank 1.

**Proposition 1.**  *$\text{Frt}(Y)$  is the free trioid of an arbitrary rank.*

If  $Y = \{x\}$ , then  $\text{Frt}(Y)$  is the free trioid of rank 1 presented by J.-L. Loday and M. O. Ronco in [1]. In the latter paper it was shown that the free associative trialgebra over a vector space is completely determined by the free associative trialgebra on one generator and the description of that trialgebra is reduced to the description of the free trioid of rank 1. A trioid which is isomorphic to the free trioid of rank 1 can be found in [4].

The notion of a normal form for elements of  $\text{Frt}(Y)$  of rank 1 (see [1], Lemma 1.10) can be naturally extended to the case of an arbitrary set  $Y$ . Namely, let  $Y$  be an arbitrary nonempty set and  $w \in \text{Frt}(Y)$ . Then we obtain the normal form for  $w$  (see [13]):

$$\begin{aligned} w &= u_1^{(0)} u_2^{(0)} \dots u_{k_0}^{(0)} \overline{u_1^{(1)}} u_2^{(1)} \dots u_{k_1}^{(1)} \overline{u_1^{(2)}} u_2^{(2)} \dots u_{k_2}^{(2)} \dots u_{k_{j-1}}^{(j-1)} \overline{u_1^{(j)}} u_2^{(j)} \dots u_{k_j}^{(j)} = \\ &= (\overline{u_1^{(0)}} \vdash \dots \vdash \overline{u_{k_0}^{(0)}}) \vdash (\overline{u_1^{(1)}} \dashv \dots \dashv \overline{u_{k_1}^{(1)}}) \perp \dots \perp (\overline{u_1^{(j)}} \dashv \dots \dashv \overline{u_{k_j}^{(j)}}), \end{aligned}$$

where  $u_l^{(i)} \in Y$ ,  $1 \leq l \leq k_i$  for all  $i \in \{0, 1, \dots, j\}$ , or

$$\begin{aligned} w &= \overline{u_1^{(1)}} u_2^{(1)} \dots u_{k_1}^{(1)} \overline{u_1^{(2)}} u_2^{(2)} \dots u_{k_2}^{(2)} \dots u_{k_{j-1}}^{(j-1)} \overline{u_1^{(j)}} u_2^{(j)} \dots u_{k_j}^{(j)} = \\ &= (\overline{u_1^{(1)}} \dashv \dots \dashv \overline{u_{k_1}^{(1)}}) \perp (\overline{u_1^{(2)}} \dashv \dots \dashv \overline{u_{k_2}^{(2)}}) \perp \dots \perp (\overline{u_1^{(j)}} \dashv \dots \dashv \overline{u_{k_j}^{(j)}}), \end{aligned}$$

where  $u_l^{(i)} \in Y$ ,  $1 \leq l \leq k_i$  for all  $i \in \{1, 2, \dots, j\}$ . Let further  $(T, \dashv', \vdash', \perp')$  be an arbitrary trioid and  $\varphi: \bar{Y} \rightarrow T$  be an arbitrary map. Since  $\text{Frt}(Y)$  is a free trioid, there exists a homomorphism  $\Phi: \text{Frt}(Y) \rightarrow (T, \dashv', \vdash', \perp')$ . It is defined by the following rule (see [13]):

$$\begin{aligned} w\Phi &= (\overline{u_1^{(0)}} \varphi \vdash' \overline{u_2^{(0)}} \varphi \vdash' \dots \vdash' \overline{u_{k_0}^{(0)}} \varphi) \vdash' (\overline{u_1^{(1)}} \varphi \dashv' \overline{u_2^{(1)}} \varphi \dashv' \dots \dashv' \overline{u_{k_1}^{(1)}} \varphi) \perp' \\ &\quad \perp' \dots \perp' (\overline{u_1^{(j)}} \varphi \dashv' \overline{u_2^{(j)}} \varphi \dashv' \dots \dashv' \overline{u_{k_j}^{(j)}} \varphi) \end{aligned}$$

or

$$\begin{aligned} w\Phi &= (\overline{u_1^{(1)}} \varphi \dashv' \overline{u_2^{(1)}} \varphi \dashv' \dots \dashv' \overline{u_{k_1}^{(1)}} \varphi) \perp' (\overline{u_1^{(2)}} \varphi \dashv' \overline{u_2^{(2)}} \varphi \dashv' \dots \dashv' \overline{u_{k_2}^{(2)}} \varphi) \perp' \\ &\quad \perp' \dots \perp' (\overline{u_1^{(j)}} \varphi \dashv' \overline{u_2^{(j)}} \varphi \dashv' \dots \dashv' \overline{u_{k_j}^{(j)}} \varphi). \end{aligned}$$

We will call  $\Phi$  the canonical homomorphism.

Now recall the definition of a dimonoid ([2, 3]).

A nonempty set  $D$  equipped with two binary associative operations  $\dashv$  and  $\vdash$  satisfying the axioms (T1) – (T3) is called a *dimonoid*. If  $D = (D, \dashv, \vdash)$  is a dimonoid, then the trioid  $(D, \dashv, \vdash, \dashv)$  (respectively,  $(D, \dashv, \vdash, \vdash)$ ) will be denoted by  $(D)^\dashv$  (respectively,  $(D)^\vdash$ ). It is clear that  $(D)^\dashv$  and  $(D)^\vdash$  are distinct as trioids but they coincide as dimonoids.

Consider some algebras from [14] and [5] which will be used in Section 4.

For an arbitrary nonempty set  $Y$  let  $Y_{\ell z} = (Y, \dashv)$ ,  $Y_{rz} = (Y, \vdash)$ ,  $Y_{rb} = Y_{\ell z} \times Y_{rz}$  be a left zero semigroup, a right zero semigroup and a rectangular band, respectively. By [14]  $Y_{\ell z, rz} = (Y, \dashv, \vdash)$  is the free left zero and right zero dimonoid (or the free left and right diband).

Define operations  $\dashv$  and  $\vdash$  on  $Y^2$  by  $(x, y) \dashv (a, b) = (x, b)$ ,  $(x, y) \vdash (a, b) = (a, b)$  for all  $(x, y), (a, b) \in Y^2$ . By [14]  $(Y^2, \dashv, \vdash)$  is the free  $(rb, rz)$ -dimonoid. It is denoted by  $Y_{rb, rz}$ .

Define operations  $\dashv$  and  $\vdash$  on  $Y^2$  by  $(x, y) \dashv (a, b) = (x, y)$ ,  $(x, y) \vdash (a, b) = (x, b)$  for all  $(x, y), (a, b) \in Y^2$ . By [14]  $(Y^2, \dashv, \vdash)$  is the free  $(\ell z, rb)$ -dimonoid. It is denoted by  $Y_{\ell z, rb}$ .

A trioid (dimonoid) is called a *rectangular triband* ([5], rectangular diband [14]), if each of its semigroups is a rectangular band.

Define operations  $\dashv$  and  $\vdash$  on  $Y^3$  by

$$(x_1, x_2, x_3) \dashv (y_1, y_2, y_3) = (x_1, x_2, y_3), \quad (x_1, x_2, x_3) \vdash (y_1, y_2, y_3) = (x_1, y_2, y_3)$$

for all  $(x_1, x_2, x_3), (y_1, y_2, y_3) \in Y^3$ . The algebra  $(Y^3, \dashv, \vdash)$  is denoted by  $\text{FRct}(Y)$ . According to Theorem 1 from [14]  $\text{FRct}(Y)$  is a free rectangular diband.

Define operations  $\dashv$ ,  $\vdash$  and  $\perp$  on  $Y^3$  by

$$(a_1, b_1, c_1) \dashv (a_2, b_2, c_2) = (a_1, b_1, c_1), \quad (a_1, b_1, c_1) \vdash (a_2, b_2, c_2) = (a_1, b_2, c_2), \\ (a_1, b_1, c_1) \perp (a_2, b_2, c_2) = (a_1, b_1, c_2)$$

for all  $(a_1, b_1, c_1), (a_2, b_2, c_2) \in Y^3$ . By Lemma 1 from [5]  $(Y^3, \dashv, \vdash, \perp)$  is a rectangular triband. It is denoted by  $Y_{\ell z, rd}$ .

Define operations  $\dashv$ ,  $\vdash$  and  $\perp$  on  $Y^3$  by

$$(a_1, b_1, c_1) \dashv (a_2, b_2, c_2) = (a_1, b_1, c_2), \quad (a_1, b_1, c_1) \vdash (a_2, b_2, c_2) = (a_2, b_2, c_2), \\ (a_1, b_1, c_1) \perp (a_2, b_2, c_2) = (a_1, b_2, c_2)$$

for all  $(a_1, b_1, c_1), (a_2, b_2, c_2) \in Y^3$ . By Lemma 2 from [5]  $(Y^3, \dashv, \vdash, \perp)$  is a rectangular triband. It is denoted by  $Y_{rd, rz}$ .

Define operations  $\dashv$ ,  $\vdash$  and  $\perp$  on  $Y^2$  by

$$(a_1, b_1) \dashv (a_2, b_2) = (a_1, b_1), \quad (a_1, b_1) \vdash (a_2, b_2) = (a_2, b_2), \quad (a_1, b_1) \perp (a_2, b_2) = (a_1, b_2)$$

for all  $(a_1, b_1), (a_2, b_2) \in Y^2$ . By Lemma 3 from [5]  $(Y^2, \dashv, \vdash, \perp)$  is a rectangular triband. It is denoted by  $Y_{\ell z, rz}^{rb}$ . Note that the trioid  $Y_{\ell z, rz}^{rb}$  was first constructed in [15].

Define operations  $\dashv$ ,  $\vdash$  and  $\perp$  on  $Y^4$  by

$$(x_1, x_2, x_3, x_4) \dashv (y_1, y_2, y_3, y_4) = (x_1, x_2, x_3, y_4), \\ (x_1, x_2, x_3, x_4) \vdash (y_1, y_2, y_3, y_4) = (x_1, y_2, y_3, y_4), \\ (x_1, x_2, x_3, x_4) \perp (y_1, y_2, y_3, y_4) = (x_1, x_2, y_3, y_4)$$

for all  $(x_1, x_2, x_3, x_4), (y_1, y_2, y_3, y_4) \in Y^4$ . The algebra  $(Y^4, \dashv, \vdash, \perp)$  is denoted by  $\text{FRT}(Y)$ . By Theorem 5 from [5]  $\text{FRT}(Y)$  is a free rectangular triband.

A nonempty subset  $A$  of a trioid  $(T, \dashv, \vdash, \perp)$  is called a subtriod, if for any  $a, b \in T$ ,  $a, b \in A$  implies  $a \dashv b, a \vdash b, a \perp b \in A$ .

As usual,  $\mathbb{N}$  denotes the set of all positive integers.

**3. Nilpotency in trioids.** In this section we introduce the notion of a nilpotent trioid, give examples of nilpotent trioids of nilpotency index 2 and construct a free  $n$ -nilpotent trioid of an arbitrary rank.

An element 0 of a trioid  $(T, \dashv, \vdash, \perp)$  is called zero ([16]), if  $x * 0 = 0 * x = 0 * 0 = 0$  for all  $x \in T$  and  $*$   $\in \{\dashv, \vdash, \perp\}$ .

A trioid  $(T, \dashv, \vdash, \perp)$  with zero will be called nilpotent, if for some  $n \in \mathbb{N}$  and any  $x_i \in T$ ,  $1 \leq i \leq n+1$ , and  $*_j \in \{\dashv, \vdash, \perp\}$ ,  $1 \leq j \leq n$ , any parenthesizing of  $x_1 *_1 x_2 *_2 \dots *_n x_{n+1}$  gives  $0 \in T$ . The least such  $n$  we shall call the nilpotency index of  $(T, \dashv, \vdash, \perp)$ . For  $k \in \mathbb{N}$  a nilpotent trioid of nilpotency index  $\leq k$  is said to be  $k$ -nilpotent.

It is clear that operations of any 1-nilpotent trioid coincide and it is a zero semigroup.

Now we give examples of nilpotent trioids of nilpotency index 2.

Let  $X_1$  and  $X_2$  be arbitrary disjoint sets,  $0 \in X_1$ , and let

$$\varphi_1: X_2 \times X_2 \rightarrow X_1, \quad \varphi_2: X_2 \times X_2 \rightarrow X_1, \quad \varphi_3: X_2 \times X_2 \rightarrow X_1$$

be arbitrary distinct maps. Define operations  $\dashv, \vdash$  and  $\perp$  on  $X_1 \cup X_2$  by

$$x \dashv y = \begin{cases} (x, y)\varphi_1, & x, y \in X_2, \\ 0, & \text{otherwise,} \end{cases} \quad x \vdash y = \begin{cases} (x, y)\varphi_2, & x, y \in X_2, \\ 0, & \text{otherwise,} \end{cases}$$

$$x \perp y = \begin{cases} (x, y)\varphi_3, & x, y \in X_2, \\ 0, & \text{otherwise} \end{cases}$$

for all  $x, y \in X_1 \cup X_2$ .

The proof of the following statement is similar to the proof of Proposition 2 from [10].

**Proposition 2.**  $(X_1 \cup X_2, \dashv, \vdash, \perp)$  is a nilpotent trioid of nilpotency index 2.

Recall that a trioid is called commutative, if its three operations are commutative.

Let  $Y$  be an arbitrary set such that  $0, a, b, c, d, e, f \in Y$  and  $a \neq b, b \neq c, c \neq d, d \neq a, b \neq e, d \neq e, f \neq e, a \neq f, c \neq f$ . Define operations  $\dashv, \vdash$  and  $\perp$  on  $Y$ , assuming

$$x \dashv y = \begin{cases} b, & x = y = a, \\ 0, & \text{otherwise,} \end{cases} \quad x \vdash y = \begin{cases} d, & x = y = c, \\ 0, & \text{otherwise,} \end{cases} \quad x \perp y = \begin{cases} f, & x = y = e, \\ 0, & \text{otherwise} \end{cases}$$

for all  $x, y \in Y$ .

The proof of the following statement is similar to that of Proposition 3 from [10].

**Proposition 3.** If  $b \neq 0$  or  $d \neq 0$ , or  $f \neq 0$ , then  $(Y, \dashv, \vdash, \perp)$  is a nilpotent commutative trioid of nilpotency index 2.

Note that the trioid  $(Y, \dashv, \vdash, \perp)$  was first constructed in [15].

It is not difficult to see that the class of all  $n$ -nilpotent trioids is a subvariety of the variety of all trioids. A trioid which is free in the variety of  $n$ -nilpotent trioids will be called a *free  $n$ -nilpotent trioid*.



See [4, 17, 18] for more information about trioids.

For every  $w \in F[X]$  denote the length of  $w$  by  $l_w$ . Let  $n \in \mathbb{N}$  and  $P_n \subset P$  be a set which contains words  $w$  with the length no more than  $n$  (see Section 2). Define operations  $\prec, \succ$  and  $\uparrow$  on the set  $P_n \cup \{0\}$  by

$$w \prec u = \begin{cases} w\tilde{u}, & l_{wu} \leq n, \\ 0, & l_{wu} > n, \end{cases} \quad w \succ u = \begin{cases} \tilde{w}u, & l_{wu} \leq n, \\ 0, & l_{wu} > n, \end{cases}$$

$$w \uparrow u = \begin{cases} wu, & l_{wu} \leq n, \\ 0, & l_{wu} > n, \end{cases} \quad w * 0 = 0 * w = 0 * 0 = 0$$

for all  $w, u \in P_n$  and  $*$   $\in \{\prec, \succ, \uparrow\}$ . Denote the algebra  $(P_n \cup \{0\}, \prec, \succ, \uparrow)$  by  $P_n^0(Y)$ .

**Theorem 1.**  $P_n^0(Y)$  is a free  $n$ -nilpotent trioid of an arbitrary rank.

*Proof.* By Proposition 1 from [16]  $P_n^0(Y)$  is a trioid with zero. For any  $w_i \in P_n^0(Y) \setminus \{0\}$ ,  $1 \leq i \leq n+1$ , and  $*_j \in \{\prec, \succ, \uparrow\}$ ,  $1 \leq j \leq n$ , any parenthesizing of  $w_1 *_1 w_2 *_2 \dots *_n w_{n+1}$  gives 0. Thus,  $P_n^0(Y)$  is nilpotent. On the other hand, for  $\bar{x} \in \bar{Y}$ ,

$$\underbrace{\bar{x} \prec \bar{x} \prec \dots \prec \bar{x}}_n = \underbrace{\bar{x}x \dots x}_n \neq 0.$$

It means that  $P_n^0(Y)$  has nilpotency index  $n$ .

Let us show that  $P_n^0(Y)$  is free in the variety of  $n$ -nilpotent trioids.

Let  $(T, \dashv', \vdash', \perp')$  be an arbitrary  $n$ -nilpotent trioid,  $\rho: \bar{Y} \rightarrow T$  be an arbitrary map and  $\mu: \text{Frt}(Y) \rightarrow (T, \dashv', \vdash', \perp')$  be the canonical homomorphism which is defined by  $\rho$  (see Section 2). Define a map  $\delta: P_n^0(Y) \rightarrow (T, \dashv', \vdash', \perp') : w \mapsto w\delta$ , assuming

$$w\delta = \begin{cases} w\mu, & w \in P_n^0(Y) \setminus \{0\}, \\ 0, & w = 0. \end{cases}$$

Show that  $\delta$  is a homomorphism.

Let  $w_1, w_2 \in P_n^0(Y) \setminus \{0\}$  and  $l_{w_1} + l_{w_2} \leq n$ . As  $w_1 \prec w_2 \in P_n^0(Y) \setminus \{0\}$ , then

$$(w_1 \prec w_2)\delta = (w_1 \prec w_2)\mu = (w_1 \dashv w_2)\mu = w_1\mu \dashv' w_2\mu = w_1\delta \dashv' w_2\delta.$$

Analogously,  $(w_1 \succ w_2)\delta = w_1\delta \vdash' w_2\delta$ ,  $(w_1 \uparrow w_2)\delta = w_1\delta \perp' w_2\delta$ . The map  $\mu$  sends an arbitrary element  $w$  to the product of some  $l_w$  elements from  $T$ . Hence, in the remaining cases the equalities

$$(w_1 \prec w_2)\delta = (w_1 \succ w_2)\delta = (w_1 \uparrow w_2)\delta = 0 = w_1\delta \dashv' w_2\delta = w_1\delta \vdash' w_2\delta = w_1\delta \perp' w_2\delta$$

hold. Thus,  $\delta$  is a homomorphism. □

**4. 0-triband decompositions of  $P_n^0(Y)$ .** In this section we introduce the notion of a 0-triband of subtrioids and in terms of 0-tribands of subtrioids describe the structure of free  $n$ -nilpotent trioids.

For trioids with zero there exists a natural analog of the notion of a triband of subtrioids (see [15]).

A trioid  $S$  with zero  $0$  (see Section 3) will be called a 0-triband of subtrioids  $S_\beta$ ,  $\beta \in B$ , where  $B$  is an idempotent trioid [15], if  $S = \bigcup_{\beta \in B} S_\beta$ ,  $S_\beta \cap S_\gamma = \{0\}$  for  $\beta \neq \gamma$  and  $S_{\beta \dashv} S_\gamma \subseteq S_{\beta+\gamma}$ ,  $S_{\beta \vdash} S_\gamma \subseteq S_{\beta+\gamma}$ ,  $S_\beta \perp S_\gamma \subseteq S_{\beta \perp \gamma}$  for any  $\beta, \gamma \in B$ . If  $B$  is an idempotent semigroup (band), then we say that  $S$  is a 0-band of subtrioids  $S_\beta$ ,  $\beta \in B$ .

Observe that the notion of a 0-triband of subtrioids generalizes the notion of a 0-diband of subdimonoids ([3]) and the notion of a 0-band of semigroups ([19]).

Let  $\omega \in F[X]$  and  $w \in P_n^0(Y) \setminus \{0\}$ . Denote the first (respectively, last) letter of  $\omega$  by  $\omega^{(0)}$  (respectively,  $\omega^{(1)}$ ). Suppose that  $u$  is the initial (respectively, terminal) subword of  $w$  with the minimal length such that  $u^{(1)} \in \bar{Y}$  (respectively,  $u^{(0)} \in \bar{Y}$ ). In this case  $\widetilde{u^{(1)}}$  (respectively,  $\widetilde{u^{(0)}}$ ) will be denoted by  $w^{[0]}$  (respectively,  $w^{[1]}$ ).

Let

$$Q_{(i,j)} = \{w \in P_n^0(Y) \setminus \{0\} \mid (\widetilde{w}^{(0)}, \widetilde{w}^{(1)}) = (i, j)\} \cup \{0\},$$

$$Q_{(i)} = \{w \in P_n^0(Y) \setminus \{0\} \mid \widetilde{w}^{(0)} = i\} \cup \{0\}, \quad Q_{[i]} = \{w \in P_n^0(Y) \setminus \{0\} \mid \widetilde{w}^{(1)} = i\} \cup \{0\}$$

for  $i, j \in Y$ ,  $n > 1$  and  $|Y| > 1$ .

The following structural theorem gives decompositions of  $P_n^0(Y)$  into 0-bands of subtrioids.

**Theorem 2.** *The free  $n$ -nilpotent trioid  $P_n^0(Y)$  is a 0-band of subtrioids*

- (i)  $Q_{(i,j)}$ ,  $(i, j) \in Y_{rb}$ , if  $n > 1$  and  $|Y| > 1$ ;
- (ii)  $Q_{(i)}$ ,  $i \in Y_{lz}$ , if  $n > 1$  and  $|Y| > 1$ ;
- (iii)  $Q_{[i]}$ ,  $i \in Y_{rz}$ , if  $n > 1$  and  $|Y| > 1$ .

*Proof.* We prove (i). It is obvious that in the case where  $n > 1$  and  $|Y| > 1$  one has  $Q_{(i,j)} \setminus \{0\} \neq \emptyset$  for all  $(i, j) \in Y_{rb}$ . Moreover,  $Q_{(i,j)}$ ,  $(i, j) \in Y_{rb}$ , is a subtrioid of  $P_n^0(Y)$ . Clearly,

$$P_n^0(Y) = \bigcup_{(i,j) \in Y_{rb}} Q_{(i,j)}, \quad Q_{(i,j)} \cap Q_{(i',j')} = \{0\}$$

for  $(i, j) \neq (i', j')$ . It is immediate to verify that

$$Q_{(i,j)} \dashv Q_{(i',j')} \subseteq Q_{(i,j')}, \quad Q_{(i,j)} \vdash Q_{(i',j')} \subseteq Q_{(i,j')}, \quad Q_{(i,j)} \perp Q_{(i',j')} \subseteq Q_{(i,j')}$$

for any  $(i, j), (i', j') \in Y_{rb}$ . So,  $P_n^0(Y)$  is a 0-band of subtrioids  $Q_{(i,j)}$ ,  $(i, j) \in Y_{rb}$ .

The proofs of the remaining cases are similar. □

Assume

$$Q_{(i,j,k,s)} = \{w \in P_n^0(Y) \setminus \{0\} \mid (\widetilde{w}^{(0)}, w^{[0]}, w^{[1]}, \widetilde{w}^{(1)}) = (i, j, k, s)\} \cup \{0\}$$

for  $i, j, k, s \in Y$ ,  $n > 3$  and  $|Y| > 1$ ;

$$Q_{(i,j,k)} = \{w \in P_n^0(Y) \setminus \{0\} \mid (\widetilde{w}^{(0)}, w^{[0]}, w^{[1]}) = (i, j, k)\} \cup \{0\},$$

$$Q_{[i,j,k]} = \{w \in P_n^0(Y) \setminus \{0\} \mid (w^{[0]}, w^{[1]}, \widetilde{w}^{(1)}) = (i, j, k)\} \cup \{0\}$$

for  $i, j, k \in Y$ ,  $n > 2$  and  $|Y| > 1$ ;  $Q_{[i,j]} = \{w \in P_n^0(Y) \setminus \{0\} \mid (w^{[0]}, w^{[1]}) = (i, j)\} \cup \{0\}$  for  $i, j \in Y$ ,  $n > 1$  and  $|Y| > 1$ .

The following two structural theorems give decompositions of  $P_n^0(Y)$  into 0-tribands of subtrioids.

**Theorem 3.** *The free  $n$ -nilpotent trioid  $P_n^0(Y)$  is a 0-triband of subtrioids*

- (i)  $Q_{(i,j,k,s)}$ ,  $(i, j, k, s) \in \text{FRT}(Y)$ , if  $n > 3$  and  $|Y| > 1$ ;
- (ii)  $Q_{(i,j,k)}$ ,  $(i, j, k) \in Y_{lz,rd}$ , if  $n > 2$  and  $|Y| > 1$ ;
- (ii)  $Q_{[i,j,k]}$ ,  $(i, j, k) \in Y_{rd,rz}$ , if  $n > 2$  and  $|Y| > 1$ ;
- (iv)  $Q_{[i,j]}$ ,  $(i, j) \in Y_{lz,rz}^{rb}$ , if  $n > 1$  and  $|Y| > 1$ .

*Proof.* We prove (i). It is easy to see that in the case where  $n > 3$  and  $|Y| > 1$  we get  $Q_{(i,j,k,s)} \setminus \{0\} \neq \emptyset$  for all  $(i, j, k, s) \in \text{FRT}(Y)$ . Furthermore,  $Q_{(i,j,k,s)}$ ,  $(i, j, k, s) \in \text{FRT}(Y)$ , is a subtrioid of  $P_n^0(Y)$ . Evidently,

$$P_n^0(Y) = \bigcup_{(i,j,k,s) \in \text{FRT}(Y)} Q_{(i,j,k,s)}, \quad Q_{(i,j,k,s)} \cap Q_{(i',j',k',s')} = \{0\}$$

for  $(i, j, k, s) \neq (i', j', k', s')$ . It can be shown that

$$\begin{aligned} Q_{(i,j,k,s)} \dashv Q_{(i',j',k',s')} &\subseteq Q_{(i,j,k,s')}, & Q_{(i,j,k,s)} \vdash Q_{(i',j',k',s')} &\subseteq Q_{(i,j',k',s')}, \\ Q_{(i,j,k,s)} \perp Q_{(i',j',k',s')} &\subseteq Q_{(i,j,k',s')} \end{aligned}$$

for any  $(i, j, k, s), (i', j', k', s') \in \text{FRT}(Y)$ . Thus,  $P_n^0(Y)$  is a 0-triband of subtrioids  $Q_{(i,j,k,s)}$ ,  $(i, j, k, s) \in \text{FRT}(Y)$ .

The proofs of the remaining cases are similar.  $\square$

Let

$$\begin{aligned} W_{(i,j,k)} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (\tilde{w}^{(0)}, w^{[0]}, \tilde{w}^{(1)}) = (i, j, k)\} \cup \{0\}, \\ W_{[i,j,k]} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (\tilde{w}^{(0)}, w^{[1]}, \tilde{w}^{(1)}) = (i, j, k)\} \cup \{0\} \end{aligned}$$

for  $i, j, k \in Y$ ,  $n > 2$  and  $|Y| > 1$ ;

$$\begin{aligned} W_{(i,j)} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (\tilde{w}^{(0)}, w^{[0]}) = (i, j)\} \cup \{0\}, \\ W_{[i,j]} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (\tilde{w}^{(0)}, w^{[1]}) = (i, j)\} \cup \{0\}, \\ W_{(i,j)} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (w^{[0]}, \tilde{w}^{(1)}) = (i, j)\} \cup \{0\}, \\ W_{[i,j]} &= \{w \in P_n^0(Y) \setminus \{0\} \mid (w^{[1]}, \tilde{w}^{(1)}) = (i, j)\} \cup \{0\}, \end{aligned}$$

$$W_{(i)} = \{w \in P_n^0(Y) \setminus \{0\} \mid w^{[0]} = i\} \cup \{0\}, \quad W_{[i]} = \{w \in P_n^0(Y) \setminus \{0\} \mid w^{[1]} = i\} \cup \{0\}$$

for  $i, j \in Y$ ,  $n > 1$  and  $|Y| > 1$ .

**Theorem 4.** *The free  $n$ -nilpotent trioid  $P_n^0(Y)$  is a 0-triband of subtrioids*

- (i)  $W_{(i,j,k)}$ ,  $(i, j, k) \in (\text{FRct}(Y))^\dagger$ , if  $n > 2$  and  $|Y| > 1$ ;
- (ii)  $W_{[i,j,k]}$ ,  $(i, j, k) \in (\text{FRct}(Y))^\dagger$ , if  $n > 2$  and  $|Y| > 1$ ;
- (iii)  $W_{(i,j)}$ ,  $(i, j) \in (Y_{lz,rb})^\dagger$ , if  $n > 1$  and  $|Y| > 1$ ;
- (iv)  $W_{[i,j]}$ ,  $(i, j) \in (Y_{lz,rb})^\dagger$ , if  $n > 1$  and  $|Y| > 1$ ;
- (v)  $W_{(i,j)}$ ,  $(i, j) \in (Y_{rb,rz})^\dagger$ , if  $n > 1$  and  $|Y| > 1$ ;
- (vi)  $W_{[i,j]}$ ,  $(i, j) \in (Y_{rb,rz})^\dagger$ , if  $n > 1$  and  $|Y| > 1$ ;

- (vii)  $W_{(i)}$ ,  $i \in (Y_{lz, rz})^\perp$ , if  $n > 1$  and  $|Y| > 1$ ;  
(viii)  $W_{[i]}$ ,  $i \in (Y_{lz, rz})^\perp$ , if  $n > 1$  and  $|Y| > 1$ .

*Proof.* We prove (i). It is readily seen that in the case where  $n > 2$  and  $|Y| > 1$  one has  $W_{(i,j,k)} \setminus \{0\} \neq \emptyset$  for all  $(i, j, k) \in (\text{FRct}(Y))^\perp$ . In addition,  $W_{(i,j,k)}$ ,  $(i, j, k) \in (\text{FRct}(Y))^\perp$ , is a subtrioid of  $P_n^0(Y)$ . It is clear that

$$P_n^0(Y) = \bigcup_{(i,j,k) \in (\text{FRct}(Y))^\perp} W_{(i,j,k)}, \quad W_{(i,j,k)} \cap W_{(i',j',k')} = \{0\}$$

for  $(i, j, k) \neq (i', j', k')$ . One can check that

$$W_{(i,j,k)} \dashv W_{(i',j',k')} \subseteq W_{(i,j,k')}, \quad W_{(i,j,k)} \vdash W_{(i',j',k')} \subseteq W_{(i,j',k')}, \quad W_{(i,j,k)} \perp W_{(i',j',k')} \subseteq W_{(i,j,k')}$$

for any  $(i, j, k), (i', j', k') \in (\text{FRct}(Y))^\perp$ . Therefore,  $P_n^0(Y)$  is a 0-triband of subtrioids  $W_{(i,j,k)}$ ,  $(i, j, k) \in (\text{FRct}(Y))^\perp$ .

The proofs of the remaining cases are similar.  $\square$

**5. The least  $n$ -nilpotent congruence on a free trioid.** In this section we present the least  $n$ -nilpotent congruence on a free trioid.

If  $f: T_1 \rightarrow T_2$  is a homomorphism of trioids, then the corresponding congruence on  $T_1$  will be denoted by  $\Delta_f$ . If  $\alpha$  is a congruence on a trioid  $(T, \dashv, \vdash, \perp)$  such that  $(T, \dashv, \vdash, \perp) / \alpha$  is an  $n$ -nilpotent trioid (see Section 3), then we say that  $\alpha$  is an  $n$ -nilpotent congruence.

Let  $\text{Frt}(Y)$  be a free trioid of an arbitrary rank (see Section 2). Fix  $n \in \mathbb{N}$  and define a relation  $\nu_n$  on  $\text{Frt}(Y)$  by

$$w_1 \nu_n w_2 \text{ if and only if } w_1 = w_2 \text{ or } l_{w_1} > n, l_{w_2} > n.$$

**Theorem 5.** *The relation  $\nu_n$  is the least  $n$ -nilpotent congruence on the free trioid  $\text{Frt}(Y)$ .*

*Proof.* Define a map  $\xi: \text{Frt}(Y) \rightarrow P_n^0(Y)$  by

$$w\xi = \begin{cases} w, & l_w \leq n, \\ 0, & l_w > n, \end{cases} \quad w \in \text{Frt}(Y).$$

Take  $w_1, w_2 \in \text{Frt}(Y)$  and assume  $l_{w_1 w_2} \leq n$ . From  $l_{w_1 w_2} \leq n$  it follows that  $l_{w_1} < n$  and  $l_{w_2} < n$ . Then

$$\begin{aligned} (w_1 \dashv w_2)\xi &= (w_1 \widetilde{w}_2)\xi = w_1 \widetilde{w}_2 = w_1 \prec w_2 = w_1 \xi \prec w_2 \xi, \\ (w_1 \vdash w_2)\xi &= (\widetilde{w}_1 w_2)\xi = \widetilde{w}_1 w_2 = w_1 \succ w_2 = w_1 \xi \succ w_2 \xi, \\ (w_1 \perp w_2)\xi &= (w_1 w_2)\xi = w_1 w_2 = w_1 \uparrow w_2 = w_1 \xi \uparrow w_2 \xi. \end{aligned}$$

If  $l_{w_1 w_2} > n$ , then

$$\begin{aligned} (w_1 \dashv w_2)\xi &= (w_1 \widetilde{w}_2)\xi = 0 = w_1 \xi \prec w_2 \xi, \\ (w_1 \vdash w_2)\xi &= (\widetilde{w}_1 w_2)\xi = 0 = w_1 \xi \succ w_2 \xi, \quad (w_1 \perp w_2)\xi = (w_1 w_2)\xi = 0 = w_1 \xi \uparrow w_2 \xi. \end{aligned}$$

Consequently,  $\xi$  is a surjective homomorphism. According to Theorem 1  $P_n^0(Y)$  is a free  $n$ -nilpotent trioid of an arbitrary rank. Then  $\Delta_\xi$  is the least  $n$ -nilpotent congruence on  $\text{Frt}(Y)$ . From the definition of  $\xi$  it follows that  $\Delta_\xi = \nu_n$ .  $\square$

## REFERENCES

1. J.-L. Loday, M.O. Ronco, *Trialgebras and families of polytopes*, Contemp. Math., **346** (2004), 369–398.
2. J.-L. Loday, *Dialgebras*, In: Dialgebras and related operads, Lect. Notes Math., Springer-Verlag, Berlin, **1763** (2001), 7–66.
3. A.V. Zhuchok, *Dimonoids*, Algebra and Logic, **50** (2011), №4, 323–340.
4. A.V. Zhuchok, *Trioids*, Asian-European Journal of Math., (2015), to appear.
5. Yul.V. Zhuchok, *Free rectangular tribands*, Buletinul Academiei de Stiinte a Republicii Moldova, Matematica, (2014), submitted.
6. A.I. Malcev, *Nilpotent semigroups*, Uchen. Zap. Ivanov. Gos. Ped. Inst., **4** (1953), 107–111. (in Russian)
7. B.H. Neumann, T. Taylor, *Subsemigroups of nilpotent groups*, Proc. Royal Soc. London, Ser. A, **274** (1963), 1–4.
8. E. Jespers, J. Okninski, *Nilpotent semigroups and semigroup algebras*, Journal of Algebra, **169** (1994), 984–1011.
9. R.S. Kruse, D.T. Price, *On the classification of nilpotent rings*, Mathematische Zeitschrift, **113** (1970), №3, 215–223.
10. A.V. Zhuchok, *Free  $n$ -nilpotent dimonoids*, Algebra and Discrete Math., **16** (2013), №2, 299–310.
11. A.V. Zhuchok, *Free  $n$ -dinilpotent dimonoids*, Problems of Physics, Mathematics and Technics, **17** (2013), №4, 43–46.
12. A.H. Clifford, *Bands of semigroups*, Proc. Amer. Math. Soc., **5** (1954), 499–504.
13. Yu.V. Zhuchok, *On definibility of free trioids by endomorphism semigroups*, Dopovidi NANU, **4** (2015), to appear. (in Russian)
14. A.V. Zhuchok, *Free rectangular dibands and free dimonoids*, Algebra and Discrete Math., **11** (2011), №2, 92–111.
15. A.V. Zhuchok, *Tribands of subtrioids*, Proc. Inst. Applied Math. and Mech., **21** (2010), 98–106.
16. A.V. Zhuchok, *On combinatorial properties of operations on trioids*, Naukovyy Chasopys NPU im. Dragomanova, Ser. 1, Phis.-Math. Nauku, **14** (2013), 77–83. (in Ukrainian)
17. A.V. Zhuchok, *Some congruences on trioids*, Journal of Mathematical Sciences, **187** (2012), №2, 138–145.
18. A.V. Zhuchok, *Semiretractions of trioids*, Ukr. Math. J., **66** (2014), №2, 218–231.
19. L.N. Shevrin, *Semigroups*, In the book: V. Artamonov, V. Salii, L. Skornyakov and others, General algebra, Sect. IV, **2** (1991), 11–191.

Department of Algebra and System Analysis  
Luhansk Taras Shevchenko National University  
yulia.mih@mail.ru

*Received 15.01.2015*